

STRING INSPIRED NEUTRINO MASS MODELS

S. F. KING

*Department of Physics and Astronomy,
University of Southampton, Southampton SO17 1BJ, U.K.
E-mail: king@soton.ac.uk*

I discuss a class of hierarchical neutrino mass models based on the see-saw mechanism with single right-handed neutrino dominance. I apply this mechanism to a string inspired SUSY Pati-Salam model and indicate how it may emerge from intersecting D-branes.

1. Introduction

Recent SNO ¹ results when combined with other solar neutrino data especially that of Super-Kamiokande strongly favour the large mixing angle (LMA) MSW solar solution ² with three active light neutrino states, and $\theta_{12} \approx \pi/6$, $\Delta m_{21}^2 \approx 5 \times 10^{-5} \text{eV}^2$ ². The atmospheric neutrino data is consistent with maximal $\nu_\mu - \nu_\tau$ neutrino mixing $\theta_{23} \approx \pi/4$ with $|\Delta m_{32}^2| \approx 2.5 \times 10^{-3} \text{eV}^2$ and the sign of Δm_{32}^2 undetermined. The CHOOZ experiment limits $\theta_{13} \lesssim 0.2$ over the favoured atmospheric range ².

The leading order Majorana mass matrix consistent with a neutrino mass hierarchy and bi-large mixing angles is

$$m_{LL} \sim \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} \frac{m}{2} \quad (1)$$

However since $m_3 \sim \sqrt{|\Delta m_{32}^2|} \sim 5.10^{-2} \text{eV}$ and $m_2 \sim \sqrt{|\Delta m_{21}^2|} \sim 7.10^{-3} \text{eV}$, compared to the natural expectation $m_2 \sim m_3$, it remains a puzzle why there should be any hierarchy at all. The question may be phrased in technical terms as one of understanding why the sub-determinant of the mass matrix is small:

$$\det \begin{pmatrix} m_{22} & m_{23} \\ m_{23} & m_{33} \end{pmatrix} \ll m^2. \quad (2)$$

2. The see-saw mechanism with single right-handed neutrino dominance

The simplest way to generate neutrino masses from a renormalisable theory is to introduce right-handed neutrinos. Since the right-handed neutrinos are electroweak singlets the Majorana masses of the right-handed neutrinos M_{RR} may be orders of magnitude larger than the Dirac masses m_{LR} . In the approximation that $M_{RR} \gg m_{LR}$ one obtains effective Majorana masses

$$m_{LL} = m_{LR} M_{RR}^{-1} m_{LR}^T \quad (3)$$

This is the see-saw mechanism^{3,4}. It not only generates Majorana mass terms of the type m_{LL} , but also naturally makes them smaller than the Dirac mass terms by a factor of $m_{LR}/M_{RR} \ll 1$. One can think of the heavy right-handed neutrinos as being integrated out to give non-renormalisable Majorana operators suppressed by the heavy mass scale M_{RR} . The goal of see-saw model building is to choose input see-saw matrices m_{LR} and M_{RR} that will give rise to Eq.1.

We now show how the input see-saw matrices can be simply chosen to give Eq.1 with a naturally small sub-determinant as in Eq.2 and hence a natural neutrino mass hierarchy using a mechanism first suggested in⁵. The idea was developed in⁶ where it was called single right-handed neutrino dominance (SRHND). SRHND was first successfully applied to the LMA MSW solution in⁷. The most recent discussion of these ideas is in⁸.

The SRHND mechanism is most simply described assuming three right-handed neutrinos in the basis where the right-handed neutrino mass matrix is diagonal although it can also be developed in other bases^{6,7}. In this basis we write the input see-saw matrices as

$$M_{RR} = \begin{pmatrix} X' & 0 & 0 \\ 0 & X & 0 \\ 0 & 0 & Y \end{pmatrix}, \quad m_{LR} = \begin{pmatrix} a' & a & d \\ b' & b & e \\ c' & c & f \end{pmatrix} \quad (4)$$

In⁵ it was suggested that one of the right-handed neutrinos may dominate the contribution to m_{LL} if it is lighter than the other right-handed neutrinos. The dominance condition was subsequently generalised to include other cases where the right-handed neutrino may be heavier than the other right-handed neutrinos but dominates due to its larger Dirac mass couplings⁶. In any case the dominant neutrino may be taken to be the third one without loss of generality. It was subsequently shown how to account for the LMA MSW solution with a large solar angle⁷ by careful consideration of the sub-dominant contributions. One of the examples considered

in ⁷ is when the right-handed neutrinos dominate sequentially,

$$\frac{|e^2|, |f^2|, |ef|}{Y} \gg \frac{|xy|}{X} \gg \frac{|x'y'|}{X'} \quad (5)$$

where $x, y \in a, b, c$ and $x', y' \in a', b', c'$. Assuming SRHND with sequential sub-dominance as in Eq.5, then Eqs.3, 4 give

$$m_{LL} \approx \begin{pmatrix} \frac{a^2}{X} + \frac{d^2}{Y} & \frac{ab}{X} + \frac{de}{Y} & \frac{ac}{X} + \frac{df}{Y} \\ . & \frac{b^2}{X} + \frac{e^2}{Y} & \frac{bc}{X} + \frac{ef}{Y} \\ . & . & \frac{c^2}{X} + \frac{f^2}{Y} \end{pmatrix} \quad (6)$$

where the contribution from the first right-handed neutrino have been neglected, but the small sub-leading contributions xy/X have been included. If the Dirac mass couplings satisfy the condition $d \ll e \approx f$ ⁵ then the matrix in Eq.6 resembles Eq.1 and furthermore has a naturally small sub-determinant as in Eq.2. The neutrino mass spectrum consists of a mass hierarchy $m_1^2 \ll m_2^2 \ll m_3^2$ where the heaviest neutrino has mass $m_3 \approx (e^2 + f^2)/Y$ ⁵. The atmospheric angle is $\tan \theta_{23} \approx e/f$ ⁵. Ignoring phases, the solar angle only depends on the sub-dominant couplings and is given by $\tan \theta_{12} \approx a/(c_{23}b - s_{23}c)$ ⁷. The simple requirement for large solar angle is then $a \sim b - c$ ⁷. Including phases the solar angle is given by an analagous result ⁸. A related phase analysis which pointed out the importance of phases for determining the solar angle was given in ⁹. One also obtains the interesting bound on the angle θ_{13} , ^{8,9,10,11} $\theta_{13} \gtrsim \frac{m_2}{m_3} \sim 0.1$. There is therefore a good chance that this angle could be observed at MINOS or CNGS. The importance of charged lepton contributions to the CHOOZ angle is explicitly discussed in ⁸.

3. A String Inspired Supersymmetric Pati-Salam Model

The motivation for considering the SUSY Pati-Salam gauge group ¹² rather than conventional SUSY GUT theories is two-fold. Firstly it avoids the infamous doublet-triplet splitting problem ¹³, and secondly it may be easier to embed the gauge group directly into a string theory ¹⁴. The problem of fermion masses in such string-inspired Pati-Salam models was first considered in ¹⁵, and these ideas were developed in detail in ^{16, 17}. The Pati-Salam gauge group ¹², supplemented by a $U(1)$ family symmetry, was first considered in ¹⁸,

$$SU(4)_{PS} \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{\text{Family}} \quad (7)$$

with left (L) and right (R) handed fermions transforming as $F_L \sim (4, 2, 1)$ and $F_R \sim (4, 1, 2)$ and the Higgs $h \sim (1, 2, 2)$ containing the two MSSM

Higgs doublets,

$$F_{L,R}^i = \begin{pmatrix} u & u & u & \nu \\ d & d & d & e^- \end{pmatrix}_{L,R}^i, \quad h = \begin{pmatrix} h_1^0 & h_2^+ \\ h_1^- & h_2^0 \end{pmatrix} \quad (8)$$

We assume the symmetry breaking sector of the minimal SUSY Pati-Salam model¹⁹ with further Higgs $H, \bar{H}$ transforming as $H \sim (4, 1, 2)$, $\bar{H} \sim (\bar{4}, 1, 2)$ and developing VEVs which break the Pati-Salam group, while $\theta, \bar{\theta}$ are Pati-Salam singlets and develop VEVs which break the $U(1)$ family symmetry. We assume for convenience that all symmetry breaking scales are at the GUT scale.

We now consider how to describe quark and lepton (including neutrino) masses and mixing angles, following²⁰. The assumed $U(1)_{\text{Family}}$ charges are

$$F_i = (1, 0, 0), \quad F_i^c = (4, 2, 0) \quad (9)$$

which leads to the universal mass matrices

$$m_{LR} \sim \begin{pmatrix} \epsilon^5 & \epsilon^3 & \epsilon \\ \epsilon^4 & \epsilon^2 & 1 \\ \epsilon^4 & \epsilon^2 & 1 \end{pmatrix}, \quad M_{RR} \sim \begin{pmatrix} \epsilon^8 & \epsilon^6 & \epsilon^4 \\ \epsilon^6 & \epsilon^4 & \epsilon^2 \\ \epsilon^4 & \epsilon^2 & 1 \end{pmatrix} \quad (10)$$

where the ϵ is some expansion parameter. The complete fermion mass operators have the form

$$(F^i F_j^c) h \left(\frac{H \bar{H}}{M^2} \right)^n \left(\frac{\theta}{M} \right)^p \quad (11)$$

where each factor of $H \bar{H}$ corresponds to a further expansion parameter power δ , with each element of the matrix having different Clebsch factors, giving vertical mass splittings within a generation. The resulting set of mass matrices, suppressing the numerical Clebsch factors and order unity coefficients, are as follows:

$$m_{LR}^\nu \sim \begin{pmatrix} \delta^3 \epsilon^5 & \delta \epsilon^3 & \delta \epsilon \\ \delta^3 \epsilon^4 & \delta^2 \epsilon^2 & \delta \\ \delta^3 \epsilon^4 & \delta^2 \epsilon^2 & 1 \end{pmatrix}, \quad m_{LR}^E \sim \begin{pmatrix} \delta \epsilon^5 & \delta \epsilon^3 & \delta \epsilon \\ \delta \epsilon^4 & \delta \epsilon^2 & \delta^2 \\ \delta \epsilon^4 & \delta \epsilon^2 & 1 \end{pmatrix} \quad (12)$$

$$m_{LR}^U \sim \begin{pmatrix} \delta^3 \epsilon^5 & \delta^2 \epsilon^3 & \delta^2 \epsilon \\ \delta^3 \epsilon^4 & \delta^2 \epsilon^2 & \delta^3 \\ \delta^3 \epsilon^4 & \delta^2 \epsilon^2 & 1 \end{pmatrix}, \quad m_{LR}^D \sim \begin{pmatrix} \delta \epsilon^5 & \delta^2 \epsilon^3 & \delta^2 \epsilon \\ \delta \epsilon^4 & \delta \epsilon^2 & \delta^2 \\ \delta \epsilon^4 & \delta \epsilon^2 & 1 \end{pmatrix} \quad (13)$$

where $\epsilon \approx \delta \approx 0.22$. An important role is played by ‘‘Clebsch zeroes’’ which forbid certain leading order operators. For example Clebsch zeroes in the 23 position forbid the leading 23 elements for the charged fermions but allow a leading 23 element for the neutrinos with a Clebsch factor 2 (not

shown explicitly). The neutrino sector has SRHND with the third right-handed neutrino dominating due to its large Dirac mass couplings which overcomes the fact that the third right-handed neutrino is the heaviest one. A large atmospheric angle is given from the large entry in the 23 position of the “lop-sided” Dirac neutrino mass matrix $\tan \theta_{23} \approx e/f \sim 1$, once the Clebsch factors and order one coefficients are included. The solar angle is $\tan \theta_{12} \sim a/(b-c) \sim 1$. The neutrino mass hierarchy is consistent with the LMA MSW solution. This model has also been shown to give successful thermal leptogenesis since the dominant right-handed neutrino is the heaviest ²¹. It also predicts a large rate for $\tau \rightarrow \mu\gamma$ partly due to large $\tan \beta \sim 50$ also due to the large 23 element of the neutrino Dirac mass matrix which leads large 23 contributions to the slepton mass matrix after renormalisation group running ²².

The above model is not based on a unified gauge group, and therefore apparently gauge unification cannot be implemented in such a model. However from the point of view of string theory the gauge couplings may be unified along with gravity at the string scale. As recently discussed ²³ the Pati-Salam model may emerge from an intersecting D-brane model as shown in Fig.1. The assumed symmetry breaking is

$$U(4)_{PS}^{(1)} \times U(4)_{PS}^{(2)} \rightarrow U(4)_{PS} \quad (14)$$

Higgs states are present which can lead to such a breaking. The $U(1)$ s are broken by the Green-Schwartz mechanism, but one $U(1)$ remains. In fact the $U(1)$ s make it hard to decouple the exotics. In the limit that the radii of compactification satisfy $R_2 \ll R_1$ we have a “single brane limit” with approximate gauge unification. The model predicts the soft mass sum rule

$$m_h^2 + m_{F_3}^2 + m_{F_3^c}^2 = M_3^2 \quad (15)$$

where m_h is the Higgs soft mass, $m_{F_3^c}$ is the right-handed third family soft mass, m_{F_3} is the left-handed third family soft mass, and M_3 is the gluino soft mass, all evaluated at the high energy scale.

References

1. N. Jelley, these proceedings.
2. A. Smirnov, these proceedings.
3. M. Gell-Mann, P. Ramond and R. Slansky in Sanibel Talk, CALT-68-709, Feb 1979, and in *Supergravity* (North Holland, Amsterdam 1979); T. Yanagida in *Proc. of the Workshop on Unified Theory and Baryon Number of the Universe*, KEK, Japan, 1979.
4. R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980) 912; J. Schechter and J. W. Valle, Phys. Rev. D **25** (1982) 774.




Figure 1. A D-brane set from which the SUSY Pati-Salam model may emerge.

5. S. F. King, Phys. Lett. B **439** (1998) 350 [arXiv:hep-ph/9806440].
6. S. F. King, Nucl. Phys. B **562** (1999) 57 [arXiv:hep-ph/9904210].
7. S. F. King, Nucl. Phys. B **576** (2000) 85 [arXiv:hep-ph/9912492].
8. S. F. King, arXiv:hep-ph/0204360.
9. S. Lavignac, I. Masina and C. A. Savoy, arXiv:hep-ph/0202086.
10. E. K. Akhmedov, G. C. Branco and M. N. Rebelo, Phys. Rev. Lett. **84** (2000) 3535 [arXiv:hep-ph/9912205].
11. F. Feruglio, A. Strumia and F. Vissani, arXiv:hep-ph/0201291.
12. J. C. Pati, A. Salam, Phys. Rev. **D10**, 275 (1974).
13. I. Antoniadis and G. K. Leontaris, Phys. Lett. B **216** (1989) 333.
14. I. Antoniadis, G. K. Leontaris and J. Rizos, Phys. Lett. B **245** (1990) 161.
15. S. F. King, Phys. Lett. B **325** (1994) 129 [Erratum-ibid. B **325** (1994) 538].
16. B. C. Allanach and S. F. King, Nucl. Phys. B **456** (1995) 57 [arXiv:hep-ph/9502219].
17. B. C. Allanach and S. F. King, Nucl. Phys. B **459** (1996) 75 [arXiv:hep-ph/9509205].
18. B. C. Allanach, S. F. King, G. K. Leontaris and S. Lola, Phys. Rev. D **56** (1997) 2632 [arXiv:hep-ph/9610517].
19. S. F. King and Q. Shafi, Phys. Lett. B **422** (1998) 135 [arXiv:hep-ph/9711288].
20. S. F. King and M. Oliveira, Phys. Rev. D **63** (2001) 095004 [arXiv:hep-ph/0009287].
21. M. Hirsch and S. F. King, Phys. Rev. D **64** (2001) 113005 [arXiv:hep-ph/0107014].
22. T. Blazek and S. F. King, Phys. Lett. B **518** (2001) 109 [arXiv:hep-ph/0105005].
23. L. L. Everett, G. L. Kane, S. F. King, S. Rigolin and L. T. Wang, Phys. Lett. B **531** (2002) 263 [arXiv:hep-ph/0202100].